

# Announcements

- Turn on the Clicker (the red LED comes on).
- Push “Join” button followed by “20” followed by the “Send” button (switches to flashing green LED if successful).
- Next exam on Chapters 13 and 14 one week from Thursday.
- Quiz on Wednesday will go through section 14.3.
- Lab reports due this week.

# Review

- Reaction Rates

- Write rate using derivative notation (Example  $A + B \rightarrow C$ ):  
$$R = -d[A]/dt = -(1/2)d[B]/dt = d[C]/dt.$$
- Rate laws of form:  $R = -d[A]/dt = k[A]^a[B]^b[C]^c \dots$
- Simple exponents (2, 1, 0, -1, -2) can easily be determined from initial rate data.
  - Rate doubles on doubling a species exponent = 1
  - Rate halves on doubling a species exponent = -1
  - Rate quadruples on doubling a species exponent = 2

- Pseudo-order (Swamping) method

- Uses large excess of all but one reactant, so concentration of only the limiting reactant (A) changes significantly.
- $-d[A]/dt = (k[B]_o^b)[A]^a \approx k_{app}[A]^a$
- For  $a = 0, 1, 2$  easily integrated to get a function for  $[A]_t$

# Simple Integrated Rate Laws

$$\text{for } -d[A]/dt = k_{\text{app}}[A]^a$$

- 0<sup>th</sup> order  $a = 0$ :  $[A]_t = [A]_o - k_{\text{app}} t$
- 1<sup>st</sup> order  $a = 1$ :  $[A]_t = [A]_o \exp\{-k_{\text{app}} t\}$ 
  - Linear:  $\ln[A]_t = \ln[A]_o - k_{\text{app}} t$
- 2<sup>nd</sup> order  $a = 2$ :  $1/[A]_t = 1/[A]_o + kt$

